

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

C A

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/535	A1	(11) International Publication Number: WO 99/37305 (43) International Publication Date: 29 July 1999 (29.07.99)
(21) International Application Number: PCT/US99/01134 (22) International Filing Date: 20 January 1999 (20.01.99) (30) Priority Data: 9801230.5 21 January 1998 (21.01.98) GB 60/072,180 22 January 1998 (22.01.98) US (71) Applicant (for all designated States except US): GLAXO GROUP LIMITED [GB/GB]; Glaxo Wellcome House, Berkeley Avenue, Greenford, Middlesex UB6 0NN (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): MORGAN, Phillip, Frederick [US/US]; Glaxo Wellcome Inc., Five Moore Drive, P.O. Box 13398, Research Triangle Park, NC 27709 (US). MUSSO, David, Lee [US/US]; Glaxo Wellcome Inc., Five Moore Drive, P.O. Box 13398, Research Triangle Park, NC 27709 (US). PARTRIDGE, John, Joseph [US/US]; Glaxo Wellcome Inc., Five Moore Drive, P.O. Box 13398, Research Triangle Park, NC 27709-3398 (US). (74) Agents: LEVY, David, J. et al.; Glaxo Wellcome Inc., Five Moore Drive, P.O. Box 13398, Research Triangle Park, NC 27709-3398 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: PHARMACEUTICALLY ACTIVE MORPHOLINOL (57) Abstract Disclosed is the compound (+)-(2S,3S)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol and pharmaceutically acceptable salts and solvates thereof, pharmaceutical compositions comprising them; also disclosed is a method of treating depression, attention deficit hyperactivity disorder (ADHD), obesity, migraine, pain, sexual dysfunction, Parkinson's disease, Alzheimer's disease, or addiction to cocaine or nicotine-containing (especially tobacco) products using such compound, salts, solvates or compositions.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.


AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LJ	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

PHARMACEUTICALLY ACTIVE MORPHOLINOL

This invention relates to an optically pure morpholinol, salts and solvates thereof, pharmaceutical formulations containing them and processes for their preparation and use.

Background of the Invention

5 Bupropion hydrochloride, (\pm)-1-(3-chlorophenyl)-2-[(1,1-dimethylethyl)-amino]-1-propanone hydrochloride, is the active ingredient of Wellbutrin® which is marketed in the United States for the treatment of depression. It is also the active ingredient of Zyban® which is marketed in the United States as an aid to smoking
10 cessation. Bupropion is a relatively weak inhibitor of the neuronal uptake of noradrenaline (NA), serotonin and dopamine (DA), and does not inhibit monoamine oxidase. While the mechanism of action of bupropion, as with other antidepressants, is unknown, it is presumed that this action is mediated by noradrenergic and/or dopaminergic mechanisms. Available evidence suggests that Wellbutrin® is a selective
15 inhibitor of noradrenaline (NA) at doses that are predictive of antidepressant activity in animal models. See Ascher, J.A., *et al.*, Bupropion: A Review of its Mechanism of Antidepressant Activity. *Journal of Clinical Psychiatry*, 56: p. 395-401, 1995.




Bupropion HCl

Bupropion is extensively metabolized in man as well as laboratory animals.
20 Urinary and plasma metabolites include biotransformation products formed via hydroxylation of the tert-butyl group and/or reduction of the carbonyl group of bupropion. Four basic metabolites have been identified. They are the erythro- and threo-amino alcohols of bupropion, the erythro-amino diol of bupropion, and a

morpholinol metabolite. These metabolites of bupropion are pharmacologically active, but their potency and toxicity relative to bupropion have not been fully characterized. Because the plasma concentrations of the metabolites are higher than those of bupropion, they may be of clinical importance.

5 The morpholinol metabolite (+/-)-(2R*,3R*)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol hydrochloride is believed to be formed from hydroxylation of the tert-butyl group of bupropion.



Morpholinol Metabolite of Bupropion HCl

In Biomed. Chromatogr. (1997), 11(3), 174-179 (Suckow, R. F *et al*) the separation and quantification of the individual enantiomers of the (+/-)-(2RS,3RS) racemic morpholinol metabolite is disclosed, as well as the fact that plasma samples contain the (-)-enantiomer to the extent of about 96% of the total of this metabolite. However, no information is given as to the pharmacological characteristics of each enantiomer and the subsequent clinical relevance thereof.


15 **Summary of the Invention**

It has now surprisingly been discovered that despite the (-) form of the morpholinol metabolite predominating significantly in human plasma samples, it is the (+) enantiomer, (+)-(2S,3S)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol in which the activity resides.

20 Thus the present invention provides, in one aspect, pharmaceutically acceptable salts and solvates of a compound of formula (I), (+)-(2S,3S)-2-(3-

chlorophenyl)-3,5,5-trimethyl-2-morpholinol.

(+)-enantiomer



5 Such pharmaceutically acceptable salts include, but are not limited to, those prepared from the following acids: hydrochloric, hydrobromic, sulfuric, nitric, phosphoric, salicylic, p-toluenesulfonic, tartaric, citric, methanesulfonic, maleic, formic, malonic, succinic, isethionic, lactobionic, naphthalene-2-sulfonic, sulfamic, ethanesulfonic and benzenesulfonic. Especially preferred is the hydrochloride salt of a
10 compound of formula (I).

Another aspect of the invention is pharmaceutical compositions comprising a compound of formula (I) or pharmaceutically acceptable salts and solvates thereof together with one or more pharmaceutically acceptable carriers, diluents or excipients.

A further aspect of the present invention is the use of a compound of formula (I)
15 or pharmaceutically acceptable salts and solvates thereof in therapy.

Yet another aspect of the invention provides methods of treating depression, attention deficit hyperactivity disorder (ADHD), obesity, migraine, pain, sexual dysfunction, Parkinson's disease, Alzheimer's disease, or addiction to cocaine or nicotine-containing (especially tobacco) products in a human or animal subject
20 comprising the administration to said subject of an effective amount of a compound of formula (I) or pharmaceutically acceptable salts and solvates thereof or pharmaceutical compositions thereof.

Yet another aspect of the present invention is the use of the compound of formula (I) or pharmaceutically acceptable salts and solvates thereof or pharmaceutical
25 compositions thereof in the preparation of a medicament for the treatment of

depression, attention deficit hyperactivity disorder (ADHD), obesity, migraine, pain, sexual dysfunction, Parkinson's disease, Alzheimer's disease, addiction to cocaine or nicotine-containing (especially tobacco) products.

5

Description of the Drawings

Figure 1. Effect of Compounds at 25 mg/kg (ip) on TBZ-Induced Depression

Figure 2. Dose Response of Compound of Formula I Against TBZ-Induced Depression (Compounds administered 30 minutes prior to TBZ, Male, CD-1 Mice, i.p., n=6)

10

Figure 3. Dose Response of Compound of Formula II Against TBZ-Induced Depression (Compounds administered 30 minutes prior to TBZ, Male, CD-1 Mice, i.p., n=6)

15

Detailed Description of the Invention

The compound of formula (I) or pharmaceutically acceptable salts and solvates thereof may be prepared by first synthesizing the racemate of the morpholinol metabolite of bupropion and subsequently separating the (+) and (-) enantiomers of the racemate via HPLC.

20

The racemate of the morpholinol metabolite of bupropion hydrochloride ((+/-)-(2R*,3R*)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol hydrochloride) may be synthesized by the following process. To 3'-chloropropiophenone (10.0g, 0.06 mol) in dioxane (50 mL) was added a solution of dioxane dibromide (14.9g, 0.06 mol) in dioxane (50 mL). The reaction mixture was stirred for 2h at ambient temperature and poured into a mixture of ice and water (500 mL). The mixture was extracted several times with methylene chloride. The combined extracts were dried (Na₂SO₄) and concentrated in vacuo to give 14.8g (85%) of 2-bromo-3'-chloropropiophenone as a pale yellow oil. This was used without further purification. NMR (300Mhz, CDCl₃); δ 7.99 (m, 1H), 7.90 (d, 1H), 7.57 (d, 1H), 7.44 (t, 1H), 5.22 (q, 1H), 1.91 (t, 3H).

25

30

To a solution of 2-bromo-3'-chloropropiophenone (19.3g, 0.08 mol) in MeOH

- (100 mL) was added dropwise a solution of 2-amino-2-methyl-1-propanol (27.8g, 0.31 mol) in methanol (200 mL) at ambient temperature. The mixture was stirred for 18h and concentrated in vacuo. The residue was partitioned between water and diethyl ether. The combined organic phase was extracted with 10% aqueous hydrogen chloride.
- 5 The combined aqueous acid extracts were chilled in an ice bath and made basic with 40% aqueous sodium hydroxide. The mixture was extracted with diethyl ether, the combined diethyl ether extracts were washed with water and saturated sodium chloride solution, dried (K_2CO_3) and concentrated in vacuo to give 15.0g (75%) of (+/-)-(2R*,3R*)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol as an off-white solid.
- 10 (+/-)-(2R*,3R*)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol may be converted to (+/-)-(2R*,3R*)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol hydrochloride by the following process. A 6.0g sample was dissolved in diethyl ether, chilled in an ice bath and ethereal hydrogen chloride added until the mixture was acidic. The resulting solid was filtered and recrystallized from ethanol/diethyl ether/ethereal
- 15 hydrogen chloride mixtures to give 4.93g of (+/-)-(2R*,3R*)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol hydrochloride as a white solid: m.p. 202-203°C. NMR (80Mhz, DMSO- d_6); δ 10.9 (br, 1H), 8.85 (br, 1H), 7.60-7.41 (m, 5H), 4.04 (d, 1H), 3.50 (d, 1H), 3.37 (br s, 1H), 1.58 (s, 3H), 1.34 (s, 3H), 1.03 (d, 3H). Anal. Calcd for $C_{13}H_{19}Cl_2NO_2$: C, 53.43; H, 6.55; N, 4.79. Found: C, 53.54; H, 6.58; N, 4.75.
- 20 (+/-)-(2R*,3R*)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol hydrochloride may be converted back to its free base by the following process. A 3.0g sample of (+/-)-(2R*,3R*)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol hydrochloride was dissolved in water (100 mL) and diethyl ether was added (200 mL). The mixture was chilled in an ice bath and the pH was adjusted to > 10 with 1.0N
- 25 aqueous sodium hydroxide. After stirring for 30 min., the phases were separated and the aqueous phase was extracted with diethyl ether. The combined diethyl ether extracts were dried (Na_2SO_4) and concentrated in vacuo to give 2.6g of (+/-)-(2R*,3R*)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol as a white solid. This was used without further purification for the chiral chromatography described below.
- 30 The (+) and (-) enantiomers of (+/-)-(2R*,3R*)-2-(3-chlorophenyl)-3,5,5-

trimethyl-2-morpholinol may be separated by the following process. (+/-)-(2R*,3R*)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol (2.54 gms.) was dissolved in 250 ml of 2 : 8 Isopropyl alcohol : Hexane (both HPLC grade). A Daicel Chiralcel OD column (2 x 25 cm.) was equilibrated for one hour at 8 ml/min. in the elution solvent, 1 : 9 : 0.2 Isopropanol : Hexane : Diethylamine. The solution of the (+/-)-(2R*,3R*)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol was injected in 1 ml. aliquots by an automated Waters Prep LC 2000, using a Waters 510 EF pump for injections. Each run was 15 minutes in length, using the conditions listed before. The separated optical isomers were collected by fraction collector (Waters) at a 2% above baseline threshold, based on 2 absorbance units full scale at 240 nm (Waters 490E UV detector). Each optical isomer solution was evaporated on a rotary evaporator at 40 degrees Centigrade and aspirator vacuum. After drying for 6 hours under high vacuum at room temperature, optical isomer 1 weighed 1.25 gm. and optical isomer 2 weighed 1.26 gm.

The enantiomeric purity of each isomer was assayed by analytical chiral HPLC on a Waters 860 HPLC with 996 Photodiode Array detector, using a Daicel Chiralcel OD-H column (4.6 x 250 mm.) eluted with 1 : 9 : 0.2 Isopropyl alcohol : Hexane : Diethylamine at 1 ml/min. Optical isomer 1 was 100% pure (R.T. 6.117 min.). Optical isomer 2 was 99.19% pure (R.T. 6.800 min.), containing 0.81% optical isomer 1 (R.T. 6.133 min.).

Hydrochloride salts of the separated enantiomers were obtained by the following processes. 1.25g (0.005 mol) of optical isomer 1 (retention time 6.117 min) ((-)-(2R,3R)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol) was dissolved in diethyl ether. The solution was filtered and the filtrate was chilled in an ice-bath adding ethereal hydrogen chloride until the solution was acidic. After standing at ambient temperature for 24h, the resulting solid was filtered, washed with diethyl ether and dried in a vacuum oven at 60° C for 18h to give 1.32g (90%) of (-)-(2R,3R)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol hydrochloride as a white solid: mp 208-209° C. NMR (300Mhz, DMSO-d₆); δ 9.72 (br, 1H), 8.76 (br, 1H), 7.54-7.41 (m, 5H), 3.98 (d, 1H), 3.52 (d, 1H), 3.37 (br s, 1H), 1.53 (s, 3H), 1.29 (s, 3H), 0.97 (d, 3H). Anal.

Calcd for $C_{13}H_{19}Cl_2NO_2$: C, 53.43; H, 6.55; N, 4.79. Found: C, 53.35; H, 6.57; N, 4.71.

$$[\alpha]_D^{20^\circ C} = -33.2^\circ (0.67, 95\% \text{ EtOH})$$

1.26g (0.005 mol) of optical isomer 2 (retention time 6.800 min) (+)-(2S,3S)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol) was dissolved in diethyl ether. The solution was filtered and the filtrate was chilled in an ice-bath adding ethereal hydrogen chloride until the solution was acidic. After standing at ambient temperature for 24h, the resulting solid was filtered, washed with diethyl ether and dried in a vacuum oven at 60° C for 18h to give 1.36g (93%) of (+)-(2S,3S)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol hydrochloride as a white solid: mp 208-209° C. NMR (300Mhz, DMSO- d_6); δ 9.87 (br, 1H), 8.76 (br, 1H), 7.54-7.41 (m, 5H), 3.99 (d, 1H), 3.51 (d, 1H), 3.37 (br s, 1H), 1.54 (s, 3H), 1.30 (s, 3H), 0.98 (d, 3H). Anal. Calcd for $C_{13}H_{19}Cl_2NO_2$: C, 53.43; H, 6.55; N, 4.79. Found: C, 53.51; H, 6.58; N, 4.73.

$$[\alpha]_D^{20^\circ C} = +31.9^\circ (0.64, 95\% \text{ EtOH})$$

The absolute configuration of (+)-(2S,3S)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol was determined by the following x-ray crystallographic method. Crystal Data: $C_{13}H_{18}Cl_2NO_2$, M=291, Orthorhombic, space group $P2_12_12_1$, $a = 8.7348$ (6), $b = 14.9824$ (10), $c = 23.1605$ (15) Å, $V=3031$ (4) Å³, $Z= 8$, $D_c= 1.276$ Mgm⁻³, $F(000)= 1226.95$. Of 12224 reflections measured. 3764 were unique and 2318 which had $I > 3.0\sigma(I)$ were used in subsequent calculations. Data was collected on a Siemens SMART diffractometer using omega scans and monochromated MoK α radiation ($\lambda = 0.71073$ Å). The positions of all non-hydrogen atoms were determined by direct methods and refined anisotropically. The hydrogen positions were all located in difference syntheses and included in subsequent refinement cycles using a riding model and an idealized bond length of 0.96 Å. The absolute configuration was determined by refinement of the Rogers' parameter and confirmed by an analysis of the 185 best Bijvoet intensity differences which indicated a probability of 0.006 that the model was in error. Least squares refinement minimized $\Sigma w(\Delta F)^2$ with weights based on counter statistics. The final agreement factors were $R_f = 0.064$ (0.108 for all data), $R_w = 0.068$ (0.081 for all data), and $GoF = 1.93$. References included E.J. Gabe, Y. Le Page, J.-P. Charland, F.L.

Lee and P.S. White, Journal of Applied Crystallography, 22, 384-387 (1989) and D. Rogers, Acta Crystallographica, A37, 734-741, 1981.

5 The amount of compound of formula (I) required to achieve the desired therapeutic effect will, of course depend on a number of factors, for example, the mode of administration, the recipient and the condition being treated. In general, the daily dose will be in the range of 0.02 to 5.0 mg/kg. More particular ranges include 0.02 to 2.5 mg/kg, 0.02 to 1.0 mg/kg, 0.02 to 0.25 mg/kg, 0.02 to 0.15 mg/kg and 0.02 to 0.07
10 mg/kg.

 The compound of formula (I) may be employed in the treatment of depression, attention deficit hyperactivity disorder (ADHD), obesity, migraine, pain, sexual dysfunction, Parkinson's disease, Alzheimer's disease, addiction to cocaine or nicotine-containing (especially tobacco) products as the compound *per se*, but is preferably
15 presented with one or more pharmaceutically acceptable carriers, diluents or excipients in the form of a pharmaceutical formulation. The carriers, diluents and excipients must, of course, be acceptable in the sense of being compatible with the other ingredients of the formulation and must not be deleterious to the recipient. The carrier may be a solid or a liquid, or both, and is preferably formulated with the agent as a unit-dose
20 formulation, for example, a tablet.

 The formulations include those suitable for oral, rectal, topical, buccal (e.g. sub-lingual) and parenteral (e.g. subcutaneous, intramuscular, intradermal or intravenous) administration.

 Formulations suitable for buccal (sub-lingual) administration include lozenges
25 comprising a compound of formula (I) in a flavoured base, usually sucrose and acacia or tragacanth, and pastilles comprising the agent in an inert base such as gelatin and glycerin or sucrose and acacia.

 Formulations of the present invention suitable for parenteral administration conveniently comprise sterile aqueous preparations of a compound of formula (I),
30 preferably isotonic with the blood of the intended recipient. These preparations are


preferably administered intravenously, although administration may also be effected by means of subcutaneous, intramuscular, or intradermal injection. Such preparations may conveniently be prepared by admixing the agent with water and rendering the resulting solution sterile and isotonic with the blood.

- 5 Formulations suitable for rectal administration are preferably presented as unit-dose suppositories. These may be prepared by admixing a compound of formula (I) with one or more conventional solid carriers, for example, cocoa butter, and then shaping the resulting mixture.

10 Formulations suitable for topical application to the skin preferably take the form of an ointment, cream, lotion, paste, gel, spray, transdermal patch, aerosol, or oil. Carriers which may be used include vaseline, lanolin, polyethylene glycols, alcohols, and combinations of two or more thereof.

15 It should be understood that in addition to the ingredients particularly mentioned above, the formulations may include other agents conventional in the art having regard to the type of formulation in question.

Biological activity of the compound of formula (I) was demonstrated by in vitro uptake models and the tetrabenazine-induced behavioural depression model. The racemic morpholinol metabolite, (+/-)-(2R*,3R*)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol, is referred to herein as "Racemate". The (-) form of the morpholinol metabolite is (-)-(2R, 3R)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol or pharmaceutically acceptable salts and solvates thereof and is referred to herein as a compound of formula (II):



In vitro Synaptosomal Uptake Experiments. In vitro uptake was determined, as reported previously, using synaptosomes prepared from rat caudoputamen (for dopamine uptake) and hypothalamus (for NA and serotonin uptake) using [³H]-dopamine, [³H]-NA and [³H]-serotonin as transport substrates, respectively. See Eckhardt, S.B., R.A. Maxwell, and R.M. Ferris, A Structure-Activity Study of the Transport Sites for the Hypothalamic and Striatal Catecholamine Uptake Systems. Similarities and differences. *Molecular Pharmacology*, 21: p. 374-9, 1982.

Synaptosomes for use in obtaining in vitro uptake data were prepared from hypothalamus or striatum by gently homogenizing the tissue in a 0.3 M sucrose/25 mM Tris pH 7.4 buffer containing iproniazid phosphate to inhibit monoamine oxidase. The homogenate was centrifuged at 1100 x g at 4°C for 10 min and the supernatant was used for uptake studies. The supernatant (~ 1 mg tissue protein) was incubated with Km concentrations of [³H]-noradrenaline, [³H]-dopamine or [³H]-serotonin at 37°C for 5 minutes in Modified Krebs-Henseleit buffer (118 mM NaCl, 5 mM KCl, 25 mM NaHCO₃, 1.2 mM NaH₂PO₄, 1.2 mM MgSO₄, 11 mM Dextrose, 2.5 mM CaCl₂) in the absence and presence of drug. Under these conditions uptake was linear with respect to both for substrate and tissue (with <5% total substrate transported). Non-specific uptake was defined as uptake at 0°C. [³H]-substrate, which had been transported into synaptosomes, was separated from free [³H]-substrate by filtration over GF/B filters and washing with cold Krebs-Henseleit buffer. The filters were counted for tritium in a liquid scintillation spectrometer.

The data for in vitro synaptosomal uptake are presented as Table 1. Among the 2 enantiomers of the morpholinol metabolite of bupropion, the (+) enantiomer, the compound of formula (I), inhibited noradrenaline (NA) uptake with an IC₅₀ of 2.2 μM. In contrast, the (-) enantiomer was ineffective at a concentration of 30 μM. On dopamine (DA) uptake, the compound of formula (I) had an IC₅₀ of ~10 μM while the (-) enantiomer was inactive at 30 μM. Neither compound inhibited serotonin uptake at 30 μM.

For comparison, Wellbutrin® was equipotent for inhibiting DA and

noradrenaline uptake with IC_{50} values of 1.9 and 2.2 μM , and did not inhibit serotonin uptake at 30 μM . Imipramine (a non-specific tricyclic antidepressant) inhibited NA uptake and serotonin uptake with IC_{50} values of 0.072 and 0.24 μM , respectively.

The compound of formula (I) was approximately twice as potent as Wellbutrin® as an NA inhibitor but, unlike the latter, was approximately 10-fold less potent as an inhibitor of dopamine uptake. These data are consistent with the observed noradrenergic actions of Wellbutrin® and the racemic morpholinol metabolite of bupropion, (+/-)-(2R*,3R*)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol hydrochloride, (306U73) in vivo, at their respective anti-TBZ doses (Cooper, B.R., *et al*, *Neuropsychopharmacology*, 11: p. 133-41,1994). Behavioral and electrophysiological data suggest that the effects of Wellbutrin® are mediated by a noradrenergic mechanism (*ibid*).

Tetrabenazine-induced Behavioural Depression Experiments. Tetrabenazine (TBZ)-induced behavioural depression was used as an in vivo measure of antidepressant activity. The test has been validated with a wide range of antidepressants, known to act through noradrenergic mechanisms (Cooper B.R. et al, "Animal models used in the prediction of antidepressant effects in man", *J. Clin. Psychiatry* 44: 63-66, 1983). Moreover, the test was also used to identify Wellbutrin® as an anti-depressant. Briefly, animals were injected with the candidate agent (p.o. or i.p.) 30 minutes before receiving an i.p. injection of tetrabenazine (35 mg/kg, as the HCl salt – prepared fresh for each use). Assessments were performed 30 minutes thereafter and included: locomotor activity (1-4 scale); ptosis (1-4 scale) and body temperature as described previously (Cooper, B.R., J.L. Howard, and F.E. Soroko, Animal models used in prediction of antidepressant effects in man (*Journal of Clinical Psychiatry*, 44: p. 63-6,1983). In all studies, the scientist performing the assessments was blind to the treatments. All parameters were weighted equally to give a "lumped" score (X) through the following algorithm:

$$X = (1 + \text{Ptosis score}) / (\text{Activity score} * [\text{Temp, treated} / \text{Temp, control}])$$

Results from the tetrabenazine-induced behavioural depression model are as follows. Assessed *in vivo* at 25 mg/kg (ip) the compound of formula (I), the racemate, Wellbutrin® and, for comparison, amitriptyline all abolished the tetrabenazine-induced behavioural depression. In contrast, the (-) enantiomer showed only modest activity (Figure 1).

In the TBZ model of behavioural depression, activity resided in the compound of formula (I). When analysed in a dose-effect study with TBZ, the activity showed a sharp increase in activity between 3 mg/kg and 6 mg/kg (ip) (Figure 2). The compound of formula II, in comparison, did not possess dose-related activity and, at 50 mg/kg, appeared to worsen the animal's condition (Figure 3). In Figures 2 and 3, AMIT (5) refers to amitriptyline dosed at 5 mg/kg and SHAM refers to a control group of animals that have received no medication at all.

Since the TBZ test has been predictive of anti-depressants acting through noradrenergic mechanisms and the compound of formula (I) is an inhibitor of noradrenaline uptake and Wellbutrin® is metabolised to this morpholinol *in vivo*, the data suggest that the anti-depressant activity of Wellbutrin® is likely to result from the effects of the compound of formula (I). (Welch, R.M., A.A. Lai, and D.H. Schroeder, Pharmacological significance of the species differences in bupropion metabolism. *Xenobiotica*, 17: p. 287-98, 1987).

By extension, other activities of Wellbutrin® could be attributed to the compound of formula (I). In particular, a noradrenergic mechanism is common to agents used to treat ADHD (e.g. methylphenidate and amphetamine). While the molecular mechanism for Wellbutrin's effects on smoking cessation is less well understood, a catecholaminergic pathway is thought to participate in the behavioural reinforcing properties of nicotine. Wellbutrin® (and, by extension, the compound of formula (I)), by augmenting NA release into brain synapses, could mimic some of the actions of nicotine and, thus, decrease the signs associated with nicotine withdrawal. Additionally, amphetamines have been used to treat obesity. The addictive properties of amphetamine, however, preclude its use for most obese patients. Wellbutrin®

causes weight loss and, like amphetamine, acts through a noradrenergic mechanism. (Zarrindast, M.R. and T. Hosseini-Nia, Anorectic and behavioural effects of bupropion. *General Pharmacology*, 19: p. 201-4,1988 and Harto-Truax, N., *et al.*, Effects of Bupropion on Body Weight. *Journal of Clinical Psychiatry*, 44: p. 183-6,1983).

5 However, unlike amphetamine, Wellbutrin® is not addictive. (Lamb, R.J. and R.R. Griffiths, Self-administration in Baboons and the Discriminative Stimulus Effects in Rats of Bupropion, Nomifensine, Diclofensine and Imipramine. *Psychopharmacology*, 102: p. 183-90,1990; Bergman, J., *et al.*, Effects of Cocaine and Related Drugs in Nonhuman Primates. III. Self-administration by Squirrel Monkeys. *Journal of*

10 *Pharmacology & Experimental Therapeutics*, 251: p. 150-5,1989 and Johanson, C.E.; and J.E. Barrett, The Discriminative Stimulus Effects of Cocaine in Pigeons. *Journal of Pharmacology & Experimental Therapeutics*, 267: p. 1-8,1993). By extension, the compound of formula (I) would also be expected to have efficacy in obesity and cocaine addiction.

15

Treatment of addiction to nicotine-containing products includes both partial and complete alleviation of addiction. Thus, in respect of tobacco products, as well as the cessation of the activity, for example smoking, this will also include reducing the level or frequency of such activity e.g. reduction of the number of cigarettes smoked in

20 a given period. In respect of other nicotine-containing products, treatment will also involve both cessation of, and a reduction in the level of, usage of such products.

Safety and Toxicity. Additional dose-ranging studies were performed to determine the range of safe doses for the isomers and the racemate. Animals were

25 observed for the presence of serious adverse events (e.g. seizures and deaths) following administration of the compounds of formula I, formula II or the racemate by the oral and intraperitoneal (i.p.) routes. The data are presented as Table II.

Administered orally, at 100 mg/kg p.o., seizures were observed with the compound of formula II and the racemate but not with the compound of formula I.

30 Seizures were observed in all of the animals with all 3 compounds when dosed at 300

mg/kg. Additionally, the 300 mg/kg oral dose resulted in 100 and 80% lethality for the compound of formula II and the racemate while no deaths were observed with the compound of formula I .

Administered i.p., all of the compounds produced seizures at 100 mg/kg. No
5 deaths were observed with the compound of formula I whereas the compound of formula II and the racemate resulted in lethality of 100% and 20 %, respectively. At the 300 mg/kg oral dose all of the lethality was observed for all of the compounds.

Table 1
Effects on Uptake *In Vitro*

[³H]-Dopamine Uptake

Compound	IC50 (mM)	SEM
Bupropion	1.9	0.15
Formula (I)	9.3	0.41
Formula (II)	>100	

[³H]-Noradrenaline Uptake

Compound	IC50 (mM)	SEM
Bupropion	2.2	0.7
Formula (I)	1.1	0.07
Formula (II)	>30	
Imipramine	0.072	0.020

[³H]-Serotonin Uptake

Compound	IC50 (mM)	SEM
Bupropion	>30	
Formula (I)	>30	
Formula (II)	>100	
Imipramine	0.24	0.03

Table 2
Adverse Events Associated with Compounds of Formula I, Formula II and the Racemate

Compound	Route	Dose (mg/kg)	Seizures (%)	Time to Seizures (min)	% Died	Time to Death (min)
Formula I	i.p.	100	100	3.93	0	n/a
Formula I	p.o.	100	0	n/a	0	n/a
Formula I	i.p.	300	100	3.95	100	6
Formula I	p.o.	300	100	11.23	0	n/a
Formula II	i.p.	100	20	5	100	7
Formula II	p.o.	100	100	7.2	0	n/a
Formula II	i.p.	300	100	1.1	100	6
Formula II	p.o.	300	100	6.8	100	7
Racemate	i.p.	100	100	3	20	14
Racemate	p.o.	100	100	9.2	0	n/a
Racemate	i.p.	300	100	3	100	3
Racemate	p.o.	300	100	6.8	80	7

N/a denotes that the effect was not observed and, therefore, no percentage was given.

Claims

1. Pharmaceutically acceptable salts and solvates of (+)-(2S,3S)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol.
- 5 2. (+)-(2S,3S)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol hydrochloride.
3. Pharmaceutical compositions comprising (+)-(2S,3S)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol or pharmaceutically acceptable salts and solvates thereof as claimed in claim 1 or claim 2 together with one or more pharmaceutically acceptable
10 carriers, diluents or excipients.
4. A compound (+)-(2S,3S)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol or a pharmaceutically acceptable salt or solvate thereof as claimed in claim 1 or claim 2 for use in therapy.
- 15 5. The use of a compound (+)-(2S,3S)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol or a pharmaceutically acceptable salt or solvate thereof as claimed in claim 1 or claim 2 or pharmaceutical compositions thereof as claimed in claim 3 in the preparation of a medicament for the treatment of depression, attention deficit
20 hyperactivity disorder (ADHD), obesity, migraine, pain, sexual dysfunction, Parkinson's disease, Alzheimer's disease, addiction to cocaine or nicotine-containing (especially tobacco) products.
- 25 6. A method of treating depression, attention deficit hyperactivity disorder (ADHD), obesity, migraine, pain, sexual dysfunction, Parkinson's disease, Alzheimer's disease, or addiction to cocaine or nicotine-containing (especially tobacco) products in a human or animal subject comprising the administration to said subject of an effective amount of (+)-(2S,3S)-2-(3-chlorophenyl)-3,5,5-trimethyl-2-morpholinol or
30 pharmaceutically acceptable salts and solvates thereof as claimed in claim 1 or claim 2 or pharmaceutical compositions thereof as claimed in claim 3.

1/3






Figure 1

2/3

**Figure 2**

3/3

**Figure 3**

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/01134

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) :A61K 31/535

US CL :514/231.2

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/231.2

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Database HCAPLUS on STN, No. 1996: 6865, Kelley, et al., "(25, 35, 5R)-2-(3, 4-Difluoro phenyl)-3, 5-dimethyl-2- morpholinol: A Novel Antidepressant Agent and Selective Inhibitor of Norepinephrine Uptake", J. Med. Chem. 39(2), pp. 347-349, 1996.	1-5
X	Database Medline on STN, No. 90341427, Martin P. et al., "Antidepressant profile of bupropion and three metabolites in mice", Pharmacopsychiatry 23(4) pp. 187-194, AZT July 1990.	1-5

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
B earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

28 APRIL 1999

Date of mailing of the international search report

03 JUN 1999

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

DWAYNE C. JONES *Lawrence Jones*

Telephone No. (703) 308-1235

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US99/01134

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

REGISTRY, HCAOLD, USPATFULL, BIOSIS, MEDLINE, EMBASE, HCAPLUS structure search with the following terms: bw306u or bw() (306u or 306), 2-(3-chloro phenyl)-2-hydroxy-3, 5, 5-trimethylmorpholine